

The replica symmetric behavior of the analogical neural network

Adriano Barra*, Giuseppe Genovese[†], Francesco Guerra[‡]

November 2009

Abstract

In this paper we continue our investigation of the analogical neural network, paying interest to its replica symmetric behavior in the absence of external fields of any type. Bridging the neural network to a bipartite spin-glass, we introduce and apply a new interpolation scheme to its free energy that naturally extends the interpolation via cavity fields or stochastic perturbations to these models.

As a result we obtain the free energy of the system as a sum rule, which, at least at the replica symmetric level, can be solved exactly. As a next step we study its related self-consistent equations for the order parameters and their rescaled fluctuations, found to diverge on the same critical line of the standard Amit-Gutfreund-Sompolinsky theory.

1 Introduction

The number of disordered models, whose description is reached in the frame of statistical mechanics for complex system, increases year by year [5]. As a consequence, the need of powerful tools for their analysis raises, which ultimately push further the global field of research suggesting new possible models where their applicability can be achieved.

Among these, interestingly, neural networks have never been analyzed from an interpolating, stochastic perturbation, perspective [18]. In fact, from the

*Dipartimento di Fisica, Sapienza Università di Roma, Piazzale Aldo Moro 2, 00185, Roma, Italy

[†]Dipartimento di Matematica, Sapienza Università di Roma, Piazzale Aldo Moro 2, 00185, Roma, Italy

[‡]Dipartimento di Fisica, Sapienza Università di Roma, and INFN, Sezione di Roma 1, Piazzale Aldo Moro 2, 00185, Roma, Italy

early work by Hopfield [27] and the, nowadays historical, theory of Amit Gutfreund and Sompolinsky (AGS) [2, 3, 4] to the modern theory for learning [9, 15], about the neural networks (thought of as spin glasses with a Hebb-like “synaptic matrix” [24]) very little is rigorously known.

Surely several contributions appeared (e.g. [1, 10, 11, 12, 13, 31, 32, 33, 34]), often following understanding of spin-glasses (e.g. [19, 20, 21, 30, 35]) and the analysis at low level of stored memories has been achieved.

However in the high level of stored memories, fundamental enquiries are still rather obscure. Furthermore general problems as the existence of a well defined thermodynamic limit, achieved for the spin glass case in [22, 23], are unsolved.

Previously we introduced an “analogical version” of the standard Hopfield model, by taking the freedom of allowing the learned patterns to live on the real axes, their probability distribution being a standard Gaussian $\mathcal{N}[0, 1]$ [7].

Within this scenario, we proved the existence of an ergodic phase where the explicit expression for all the thermodynamical quantities (free energy, entropy, internal energy) have been found to self-average around their annealed expression in the thermodynamic limit, in complete agreement with AGS theory.

In this paper, again by using an analogy among neural networks and bipartite spin glasses, we move on introducing a novel interpolating technique (essentially based on two different stochastic perturbations) which we use to give a complete description of the analogical Hopfield model phase diagram in the replica symmetric approximation and with high level of stored memories (i.e. patterns).

Furthermore we control the fluctuations and correlations of the order parameters of the theory, whose divergences confirm the transition line predicted by standard AGS theory to hold even in this continuous counterpart.

As a last remark we stress that the whole is exploited without external fields checking system responses and, as a consequence, nor retrieval neither the presence of any “magnetization” are discussed and are left for future speculation.

The paper is organized as follows: In Sec. 2 we introduce the analogical neural network with all its statistical mechanics package of definitions and properties. In Sec. 3 we analyze its replica symmetric behavior by means of our interpolating scheme, while in Sec. 4 we exploit the fluctuation control to check for regularities and singularities of the order parameters, obtaining the critical line for the phase transition from the ergodic regime to a non-ergodic one.

Sec. 5 is left for conclusion and outlook.

2 Analogical neural network

We introduce a large network of N two-state neurons $\sigma_i = \pm 1$, $i \in (1, \dots, N)$, which are thought of as quiescent (sleeping) when their value is -1 or spiking (emitting a current signal to other neurons) when their value is $+1$. They interact throughout a symmetric synaptic matrix J_{ij} defined accordingly the Hebb rule for learning,

$$J_{ij} = \sum_{\mu=1}^k \xi_i^\mu \xi_j^\mu. \quad (1)$$

Each random variable $\xi^\mu = \{\xi_1^\mu, \dots, \xi_N^\mu\}$ represents a learned pattern and tries to bring the overall current in the network (or in some part) stable with respect to itself (when this happens, we say we have a retrieval state, see e.g. [2]). The analysis of the network assumes that the system has already stored p patterns (no learning is investigated) and we are interested in the case in which this number increases proportionally (linearly) to the system size (high storage level).

In standard literature these patterns are usually taken at random with distribution $P(\xi_i^\mu) = (1/2)\delta_{\xi_i^\mu, +1} + (1/2)\delta_{\xi_i^\mu, -1}$, while we extend their support to be on the real axes weighted by a Gaussian probability distribution, i.e.

$$P(\xi_i^\mu) = \frac{1}{\sqrt{2\pi}} e^{-(\xi_i^\mu)^2/2}. \quad (2)$$

The Hamiltonian of the model is defined as follows

$$H_N(\sigma; \xi) = -\frac{1}{N} \sum_{\mu=1}^k \sum_{i < j}^N \xi_i^\mu \xi_j^\mu \sigma_i \sigma_j, \quad (3)$$

which, splitting the summations $\sum_{i < j}^N = \frac{1}{2} \sum_{ij}^N - \frac{1}{2} \sum_i^N \delta_{ij}$ enable us to write down the following partition function

$$\begin{aligned} Z_N(\beta; \xi) &= \sum_{\sigma} \exp \left(\frac{\beta}{2N} \sum_{\mu=1}^k \sum_{ij}^N \xi_i^\mu \xi_j^\mu \sigma_i \sigma_j - \frac{\beta}{2N} \sum_{\mu=1}^k \sum_i^N (\xi_i^\mu)^2 \right) \quad (4) \\ &= \tilde{Z}(\beta; \xi) \times \left(e^{\frac{-\beta}{2N} \sum_{\mu=1}^k \sum_{i=1}^N (\xi_i^\mu)^2} \right). \end{aligned}$$

β , the inverse temperature in spin glass theory, denotes the level of noise in the network and we defined

$$\tilde{Z}(\beta; \xi) = \sum_{\sigma} \exp\left(\frac{\beta}{2N} \sum_{\mu=1}^k \sum_{ij}^N \xi_i^{\mu} \xi_j^{\mu} \sigma_i \sigma_j\right). \quad (5)$$

Notice that the last term at the r.h.s. of eq. (4) does not depend on the particular state of the network.

As a consequence, the control of the last term easily follows:

$$\ln Z_{N,k}(\beta; \xi) = \ln \tilde{Z}_{N,k}(\beta; \xi) - \frac{\beta}{2N} \sum_{\mu}^k \sum_i^N (\xi_i^{\mu})^2 = \ln \tilde{Z}_{N,k}(\beta; \xi) - \frac{\beta}{2} \hat{f}_N \quad (6)$$

where, as $\hat{f}_N$ is a sum of independent random variables, $\mathbb{E} \hat{f}_N = k$ and $\lim_{N \rightarrow \infty} (1/N) \mathbb{E} \hat{f}_N = k/N$, which in the thermodynamic limit, simply adds a term $-\alpha\beta/2$ to the free energy (to be defined in (11)).

Consequently we focus just on $\tilde{Z}(\beta; \xi)$. Let us apply the Gaussian integration [16] to linearize with respect to the bilinear quenched memories carried by the $\xi_i^{\mu} \xi_j^{\mu}$: The expression for the partition function (5) becomes (renaming $\tilde{Z} \rightarrow Z$ for simplicity)

$$Z_N(\beta; \xi) = \sum_{\sigma} \int \prod_{\mu=1}^k d\mu(z_{\mu}) \exp\left(\sqrt{\frac{\beta}{N}} \sum_{\mu=1}^k \sum_{i=1}^N \xi_i^{\mu} \sigma_i z_{\mu}\right), \quad (7)$$

with $d\mu(z_{\mu})$ standard Gaussian measure for all the z_{μ} .

Taken O as a generic function of the neurons, we define the Boltzmann state $\omega_{\beta}(O)$ at a given level of noise β as

$$\omega_{\beta}(O) = \omega(O) = (Z_N(\beta; \xi))^{-1} \sum_{\sigma} O(\sigma) e^{-\beta H_N(\sigma; \xi)}, \quad (8)$$

and often we drop the subscript β for the sake of simplicity. The s -replicated Boltzmann measure is defined as $\Omega = \omega^1 \times \omega^2 \times \dots \times \omega^s$ in which all the single Boltzmann states are independent states at the same noise level β^{-1} and share an identical distribution of quenched memories ξ . For the sake of clearness, given a function F of the neurons of the s replicas and the freedom of using the symbol $a \in [1, \dots, s]$ to label replicas, such an average can be written as

$$\Omega(F(\sigma^1, \dots, \sigma^s)) = \frac{1}{Z_N^s} \sum_{\sigma^1} \sum_{\sigma^2} \dots \sum_{\sigma^s} F(\sigma^1, \dots, \sigma^s) \exp\left(-\beta \sum_{a=1}^s H_N(\sigma^a, \xi)\right). \quad (9)$$

The average over the quenched memories will be denoted by $\mathbb{E}$ and for a generic function of these memories $F(\xi)$ can be written as

$$\mathbb{E}[F(\xi)] = \int \prod_{\mu=1}^p \prod_{i=1}^N \frac{d\xi_i^\mu e^{-\frac{(\xi_i^\mu)^2}{2}}}{\sqrt{2\pi}} F(\xi) = \int F(\xi) d\mu(\xi), \quad (10)$$

of course $\mathbb{E}[\xi_i^\mu] = 0$ and $\mathbb{E}[(\xi_i^\mu)^2] = 1$.

We use the symbol $\langle . \rangle$ to mean $\langle . \rangle = \mathbb{E}\Omega(.).$

In the thermodynamic limit, it is assumed

$$\lim_{N \rightarrow \infty} \frac{p}{N} = \alpha,$$

α being a given real number, parameter of the theory.

For the sake of simplicity we allow a little abuse in the notation so to use the symbol α even at finite N , still meaning the ration among the two parties.

The main quantity of interest is the quenched intensive pressure defined as

$$A_N(\alpha, \beta) = -\beta f_N(\alpha, \beta) = \frac{1}{N} \mathbb{E} \ln Z_N(\beta; \xi). \quad (11)$$

Here, $f_N(\alpha, \beta) = u_N(\alpha, \beta) - \beta^{-1} s_N(\alpha, \beta)$ is the free energy density, $u_N(\alpha, \beta)$ the internal energy density and $s_N(\alpha, \beta)$ the intensive entropy.

Reflecting the bipartite nature of the Hopfield model expressed by eq. (7) we introduce two other order parameters: the first is the overlap between the replicated neurons (first party overlap), defined as

$$q_{ab} = \frac{1}{N} \sum_{i=1}^N \sigma_i^a \sigma_i^b \in [-1, +1], \quad (12)$$

and the second the overlap between the replicated Gaussian variables z (second party overlap), defined as

$$p_{ab} = \frac{1}{p} \sum_{\mu=1}^k z_a^\mu z_b^\mu \in (-\infty, +\infty). \quad (13)$$

Both the two order parameters above play a considerable role in the theory as they can express thermodynamical quantities [7].

3 Replica symmetric free energy

In this section we pay attention to the structure of the free energy: we want to obtain the latter via a sum rule in which we may isolate explicitly the order parameter fluctuations so to be able to neglect them achieving a replica-symmetric behavior.

Due to the equivalence among neural network and bipartite spin-glasses, we generalize the way cavity field and the stochastic stability techniques work on spin glasses to these structures by introducing a new interpolation scheme as follows:

For the sake of clearness, in order to exploit the interpolation method adapted to the physics of the model, we introduce 3 free parameters in the interpolating structure (i.e. a, b, c) that we fix a fortiori, once the sum rule is almost achieved.

In a pure stochastic stability fashion [20], we need to introduce also two classes of i.i.d. $\mathcal{N}[0, 1]$ variables, namely N variables η_i and K variables $\tilde{\eta}_\mu$, whose average is still encoded into the $\mathbb{E}$ operator and by which we define the following interpolating quenched pressure $\tilde{A}_{N,k}(\beta, t)$

$$\begin{aligned} \tilde{A}_{N,k}(\beta, t) = & \frac{1}{N} \mathbb{E} \log \sum_{\sigma} \int \prod_{\mu} d\mu(z_{\mu}) \exp(\sqrt{t} \frac{\beta}{N} \sum_{i,\mu}^{N,k} \xi_i^{\mu} \sigma_i z_{\mu}) \quad (14) \\ & \cdot \exp(a\sqrt{1-t} \sum_i^N \eta_i \sigma_i) \exp(b\sqrt{1-t} \sum_{\mu}^k \tilde{\eta}_{\mu} z_{\mu}) \exp(c \frac{(1-t)}{2} \sum_{\mu}^k z_{\mu}^2). \end{aligned}$$

We stress that $t \in [0, 1]$ interpolates between $t = 0$ where the interpolating quenched pressure becomes made of by non-interacting systems (a series of one-body problem) whose integration is straightforward and the opposite limit, $t = 1$, that recovers the correct quenched free energy (11).

The plan is then to evaluate the t -streaming of such a quantity and than obtain the correct expression by using the fundamental theorem of calculus:

$$A_{N,k}(\beta) = \tilde{A}_{N,k}(\beta, t = 1) = \tilde{A}_{N,k}(\beta, t = 0) + \int_0^1 dt' \left(\partial_{t'} \tilde{A}_{N,k}(\beta, t') \right)_{t=t'}. \quad (15)$$

When evaluating the streaming $\partial_t \tilde{A}$ we get the sum of four terms (A, B, C, D): each comes as a consequence of the derivation of a corresponding exponential term appearing into the expression (14).

Once introduced the averages $\langle \cdot \rangle_t$ that naturally extend the Boltzmann measure encoded in the interpolating scheme (and reduce to the proper one

whenever setting $t = 1$), we can write them down as

$$\begin{aligned}
A &= \frac{1}{N} \sqrt{\frac{\beta}{N}} \frac{1}{2\sqrt{t}} \sum_{i,\mu}^{N,k} \mathbb{E} \xi_{i,\mu} \omega(\sigma_i z_\mu) = \frac{\beta}{2N} \mathbb{E} \sum_{\mu}^k \omega(z_\mu^2) - \frac{\alpha\beta}{2} \langle q_{12} p_{12} \rangle_t, \\
B &= \frac{-a}{2N\sqrt{1-t}} \sum_i^N \mathbb{E} \eta_i \omega(\sigma_i) = -\frac{a^2}{2} (1 - \langle q_{12} \rangle_t), \\
C &= \frac{-b}{2N\sqrt{1-t}} \sum_{\mu}^k \mathbb{E} \tilde{\eta}_{\mu} \omega(z_{\mu}) = \frac{-b^2}{2N} \sum_{\mu}^k \mathbb{E} \omega(z_{\mu}^2) + \frac{\alpha b^2}{2} \langle p_{12} \rangle_t, \\
D &= \frac{-c}{2N} \sum_{\mu}^k \omega(z_{\mu}^2),
\end{aligned}$$

where in the first three equations we used integration by parts (Wick theorem).

In the replica symmetric ansatz, the order parameters do not fluctuate with respect to the quenched average and the only values (at any given β, α point) they gets are $\langle q \rangle = \bar{q}$, $\langle p \rangle = \bar{p}$, where the bars denote the replica symmetric approximation.

Summing all the contributions (A, B, C, D) and adding and subtracting the term $\alpha\beta\bar{q}\bar{p}/2$ (that we use to center and complete the square of the two overlaps), we get

$$\begin{aligned}
\frac{d\tilde{A}_{N,k}(\beta, t)}{dt} &= (\beta - b^2 - c) \frac{1}{2N} \mathbb{E} \sum_{\mu}^k \omega(z_{\mu}^2) - \frac{\alpha\beta}{2} \langle q_{12} p_{12} \rangle_t - \\
&- \frac{a^2}{2} (1 - \langle q_{12} \rangle_t) + \frac{\alpha b^2}{2} \langle p_{12} \rangle_t + \frac{\alpha\beta}{2} \bar{q}\bar{p} - \frac{\alpha\beta}{2} \bar{q}\bar{p}.
\end{aligned} \tag{16}$$

So we see that if we choose

$$a = \sqrt{\alpha\beta\bar{p}}, \quad b = \sqrt{\beta\bar{q}}, \quad c = \beta(1 - \bar{q}),$$

we get

$$\frac{d\tilde{A}_{N,k}(\beta, t)}{dt} = -\frac{\alpha\beta}{2} \langle (q_{12} - \bar{q})(p_{12} - \bar{p}) \rangle_t - \frac{\alpha\beta}{2} \bar{p}(1 - \bar{q}). \tag{17}$$

Once inserted the expression (17) into eq.(15) the sum rule for the free energy is achieved.

In order to get the replica symmetric solution $A_{N,k}^{RS}(\beta)$ we impose the self-averaging of the overlaps, so that we need to evaluate only

$$A_{N,k}^{RS}(\beta) = \tilde{A}_{N,k}(\beta, t=0) - \frac{\alpha\beta}{2}\bar{p}(1-\bar{q}) - \frac{\alpha\beta}{2}, \quad (18)$$

where the last term at the r.h.s. comes from the diagonal term of the first party as explained in Sec. 2.

The evaluation of $\tilde{A}_{N,k}(\beta, t=0)$ is easily available because it is a one-body calculation, which implies factorization in the volume sizes. Namely, we have to evaluate explicitly the quantity

$$\begin{aligned} \tilde{A}_{N,k}(\beta, t=0) &= \quad (19) \\ &= \frac{1}{N} \mathbb{E} \log \sum_{\sigma} \int \prod_{\mu}^k d\mu(z_{\mu}) e^{\sqrt{\alpha\beta\bar{p}} \sum_i^N \eta_i \sigma_i} e^{\sqrt{\beta\bar{q}} \sum_{\mu}^k \tilde{\eta}_{\mu} z_{\mu}} e^{\frac{\beta}{2}(1-\bar{q}) \sum_{\mu}^p z_{\mu}^2} \\ &= \frac{1}{N} \mathbb{E} \log \sum_{\sigma} e^{\sqrt{\alpha\beta\bar{p}} \sum_i^N \eta_i \sigma_i} + \\ &\quad + \frac{1}{N} \mathbb{E} \log \int \prod_{\mu}^k dz_{\mu} e^{-\frac{1}{2} \sum_{\mu}^k z_{\mu}^2 (1-\beta(1-\bar{q}))} e^{\sqrt{\beta\bar{q}} \sum_{\mu}^k \eta_{\mu} z_{\mu}} \\ &= \log 2 + \int d\mu(\eta) \log \cosh(\sqrt{\alpha\beta\bar{p}}\eta) + \\ &\quad + \frac{\alpha}{2} \log(1 - \beta(1 - \bar{q})) + \alpha \mathbb{E} \log \int dr e^{-r^2/2} e^{\sqrt{\frac{\beta\bar{q}}{1-\beta(1-\bar{q})}} \eta r}, \end{aligned}$$

where we introduced $r = \sigma z$, σ defining the standard Gaussian variance such that

$$\sigma^2 = (1 - \beta(1 - \bar{q}))^{-1}. \quad (20)$$

As a consequence we get

$$\begin{aligned} \tilde{A}_{N,k}(\beta, t=0) &= \log 2 + \int d\mu(\eta) \log \cosh(\sqrt{\alpha\beta\bar{p}}\eta) + \quad (21) \\ &\quad + \frac{\alpha}{2} \log\left(\frac{1}{1 - \beta(1 - \bar{q})}\right) + \frac{\alpha\beta}{2} \frac{\bar{q}}{1 - \beta(1 - \bar{q})}, \end{aligned}$$

and, overall, we can state the next

Theorem 1. *The replica symmetric free energy of the analogical Hopfield*

neural network is given by the following expression

$$A^{RS}(\beta, \alpha) = \log 2 + \int d\mu(\eta) \log \cosh(\sqrt{\alpha\beta\bar{p}}\eta) + \quad (22)$$

$$+ \frac{\alpha}{2} \log\left(\frac{1}{1 - \beta(1 - \bar{q})}\right) + \frac{\alpha\beta}{2} \frac{\bar{q}}{1 - \beta(1 - \bar{q})} - \frac{\alpha\beta}{2} \bar{p}(1 - \bar{q}) - \frac{\alpha\beta}{2}.$$

Remark 1. We stress that in the ergodic regime, where the overlap self-averages to zero, the expression recover the correct ergodic expression [7] as well as the annealed expression of the Sherrington-Kirkpatrick model (SK) when sending $\alpha \rightarrow \infty$ and $\beta \rightarrow 0$ by keeping $\alpha\beta = \beta_{SK}$.

Self-consistency relations can be found by imposing equal to zero the partial derivatives of the free energy with respect to its order parameters, namely the system $(\partial_q A(\beta, \alpha) = 0), (\partial_p A(\beta, \alpha) = 0)$, which gives

$$\frac{\partial A}{\partial q} = \frac{\alpha\beta}{2} \left(\bar{p} - \frac{\beta\bar{q}}{(1 - \beta(1 - q))^2} \right) = 0 \quad (23)$$

$$\frac{\partial A}{\partial p} = \frac{\alpha\beta}{2} \left(\int d\mu(z) \tanh^2(\sqrt{\alpha\beta\bar{p}}z) - \bar{q} \right) = 0, \quad (24)$$

by which

$$\bar{q} = \int d\mu(z) \tanh^2 \left(\frac{\sqrt{\alpha\bar{q}}\beta z_\mu}{(1 - \beta(1 - \bar{q}))} \right), \quad (25)$$

and as a consequence $\bar{p}(\bar{q}) = \beta\sigma^4\bar{q}$. These conditions can be seen as a minimax principle defining the replica symmetric solution. Let us recall that in the spin glass case we have a minimum principle instead [21].

4 Fluctuations of the order parameters and critical line

We are now ready to separate different regions in the phase diagram, where different behaviors do appear. In particular we want to see where the annealing, characterized by $(q = 0, p = 0)$, is spontaneously broken and ergodicity is lost.

To satisfy this task we proceed as follows: at first we introduce the streaming equation so to be able to calculate variations of generic observable as overlap correlation functions.

Then we define the centered and rescaled overlaps and introduce their correlation matrix. Each element of this matrix then is evaluated at $t = 0$

and then propagated thought $t = 1$ via its streaming: This procedure encodes naturally for a system of coupled linear differential equations that, once solved, give the expressions of the overlap fluctuations. The latter are found to diverge on a line in the (α, β) plane, which becomes a natural candidate for a second order phase transition (confirmed by the regularity of the behavior before such a line is reached from the ergodic phase). Let us start the plan by introducing the following

Proposition 1. *Given O as a smooth function of s replica overlaps $(q_1, \dots, q_s)$ and $(p_1, \dots, p_s)$, the following streaming equation holds:*

$$\begin{aligned} \frac{d}{dt}\langle O \rangle_t &= \beta\sqrt{\alpha} \left(\sum_{a,b}^s \langle O \cdot \xi_{a,b} \eta_{a,b} \rangle_t \right. \\ &\quad \left. - s \sum_{a=1}^s \langle O \cdot \xi_{a,s+1} \eta_{a,s+1} \rangle_t + \frac{s(s+1)}{2} \langle O \cdot \xi_{s+1,s+2} \eta_{s+1,s+2} \rangle_t \right). \end{aligned} \quad (26)$$

We skip the proof as is long but simple and works by direct evaluation pretty standard in the disordered system literature (see for example [21, 6, 8]).

The rescaled overlap ξ_{12} and η_{12} are defined accordingly to

$$\xi_{12} = \sqrt{N} (q_{12} - \bar{q}), \quad (27)$$

$$\eta_{12} = \sqrt{K} (p_{12} - \bar{p}). \quad (28)$$

In order to control the overlap fluctuations, namely $\langle \xi_{12}^2 \rangle_{t=1}$, $\langle \xi_{12} \eta_{12} \rangle_{t=1}$, $\langle \eta_{12}^2 \rangle_{t=1}$, ..., noting that the streaming equation pastes two replicas to the ones already involved ($s = 2$ so far), we need to study nine correlation functions. It is then useful to introduce them and link them to capital letters so to simplify their visualization:

$$\langle \xi_{12}^2 \rangle_t = A(t), \quad \langle \xi_{12} \xi_{13} \rangle_t = B(t), \quad \langle \xi_{12} \xi_{34} \rangle_t = C(t), \quad (29)$$

$$\langle \xi_{12} \eta_{12} \rangle_t = D(t), \quad \langle \xi_{12} \eta_{13} \rangle_t = E(t), \quad \langle \xi_{12} \eta_{34} \rangle_t = F(t), \quad (30)$$

$$\langle \eta_{12} \eta_{12} \rangle_t = G(t), \quad \langle \eta_{12} \eta_{13} \rangle_t = H(t), \quad \langle \eta_{12} \eta_{34} \rangle_t = I(t). \quad (31)$$

Let us now sketch their streaming. Let us at first introduce the operator “dot” as

$$\dot{O} = \frac{1}{\beta\sqrt{\alpha}} \frac{dO}{dt},$$

which simplifies calculations and shifts the propagation of the streaming from $t = 1$ to $t = \beta\sqrt{\alpha}$: Using it we sketch how to write the streaming of the first two correlations (as it works in the same way for any other):

$$\begin{aligned}\dot{A} &= \langle \xi_{12}^2 \xi_{12} \eta_{12} \rangle_t - 4 \langle \xi_{12}^2 \xi_{13} \eta_{13} \rangle_t + 3 \langle \xi_{12}^2 \xi_{34} \eta_{34} \rangle_t, \\ \dot{B} &= \langle \xi_{12} \xi_{13} (\xi_{12} \eta_{12} + \xi_{13} \eta_{13} + \xi_{23} \eta_{23}) \rangle_t - \\ &\quad - 3 \langle \xi_{12} \eta_{13} (\xi_{14} \eta_{14} + \xi_{24} \eta_{24} + \xi_{34} \eta_{34}) \rangle_t + 6 \langle \xi_{12} \eta_{13} \xi_{45} \eta_{45} \rangle_t.\end{aligned}$$

By assuming a Gaussian behavior, as in the strategy outlined in [21], we can write the overall streaming of the correlation functions in the form of the following differential system

$$\begin{aligned}\dot{A} &= 2AD - 8BE + 6CF, \\ \dot{B} &= 2AE + 2BD - 4BE - 6BF - 6EC + 12CF, \\ \dot{C} &= 2AF + 2CD + 8BE - 16BF - 16CE + 20CF, \\ \dot{D} &= AG - 4BH + 3CI + D^2 - 4E^2 + 3F^2, \\ \dot{E} &= AH + BG - 2BH - 3BI - 3CH + 6CI + 2ED - 2E^2 - 6EF + 6F^2, \\ \dot{F} &= AI + CG + 4BH - 8BI - 8CH + 10CI + 2DF + 4E^2 - 16EF + 10F^2, \\ \dot{G} &= 2GD - 8HE + 6IF, \\ \dot{H} &= 2GE + 2HD - 4HE - 6HF - 6IE + 12IF, \\ \dot{I} &= 2GF + 2DI + 8HE - 16HF - 16IE + 20IF.\end{aligned}$$

It is easy to solve this system, once the initial conditions at $t = 0$ are known. Our general analysis covers also the case where external fields are involved. We do not report here the full analysis, for the sake of brevity.

In order to proceed further, in our case of absence of external fields, we need to evaluate these correlations at $t = 0$. As at $t = 0$ everything is factorized, the only needed check is by the correlations inside each party.

Starting with the first party, we have to study A, B, C at $t = 0$. As only the diagonal terms give not negligible contribution, it is immediate to work out this first set of starting points as

$$A(0) = N^{-1} \sum_i^N (1 - 2\bar{q} \langle \sigma_i^1 \sigma_i^2 \rangle + \bar{q}^2) = 1 - \bar{q}^2, \quad (32)$$

$$B(0) = N^{-1} \sum_i^N (\sigma_i^2 \sigma_i^3 - \bar{q} \sigma_i^1 \sigma_i^2 - \bar{q} \sigma_i^1 \sigma_i^3 + \bar{q}^2) = \bar{q} - \bar{q}^2, \quad (33)$$

$$\begin{aligned} C(0) &= N^{-1} \sum_{ij}^{N,N} (\sigma_i^1 \sigma_i^2 \sigma_i^3 \sigma_i^4 - \bar{q} \sigma_i^1 \sigma_i^2 - \bar{q} \sigma_j^3 \sigma_j^4 + \bar{q}^2) = \\ &= \int d\mu(z) \tanh^4\left(\frac{\beta \sqrt{\alpha \bar{q}} z}{1 - \beta(1 - \bar{q})}\right) - \bar{q}^2, \end{aligned} \quad (34)$$

where we stress that even in the last equation only the diagonal terms $i = j$ contribute.

For the second party we need to evaluate G, H, I at $t = 0$. The only difference with the first party is the lacking of the dichotomy of its elements such that $z_\mu^2 \neq 1$ as for the σ 's.

It is immediate to check that $G(0), H(0), I(0)$ are function of $\omega(z^2)$ and $\omega^2(z)$, which are Gaussian integrals and can be worked out as

$$\omega(z) = \frac{\int z e^{\sqrt{\beta \bar{q}} \eta z} e^{\frac{\beta}{2}(1-\bar{q})z^2} e^{-z^2/2} dz}{\int e^{\sqrt{\beta \bar{q}} \eta z} e^{\frac{\beta}{2}(1-\bar{q})z^2} e^{-z^2/2} dz} = \sqrt{\beta \bar{q}} \eta \sigma^2, \quad (35)$$

$$\omega(z^2) = \frac{\int z e^{\sqrt{\beta \bar{q}} \eta z} e^{\frac{\beta}{2}(1-\bar{q})z^2} e^{-z^2/2} dz}{\int e^{\sqrt{\beta \bar{q}} \eta z} e^{\frac{\beta}{2}(1-\bar{q})z^2} e^{-z^2/2} dz} = \sigma^2 (1 + \sqrt{\beta \bar{q}} \eta \sigma)^2. \quad (36)$$

Remembering that $\beta \sigma^4 \bar{q} = \bar{p}$ (cfr. eq.(24)), we get

$$\begin{aligned} G(0) &= \mathbb{E} \omega(z^2) \omega(z^2) - \bar{p}^2 = \mathbb{E} \sigma^4 (1 + \sqrt{\beta \bar{q}} \eta \sigma)^4 - \bar{p}^2, \\ H(0) &= \mathbb{E} \omega(z^2) \omega(z)^2 - \bar{p}^2 = \mathbb{E} \sigma^2 (1 + \sqrt{\beta \bar{q}} \eta \sigma)^2 \beta \bar{q} \eta^2 \sigma^4 - \bar{p}^2, \\ I(0) &= \mathbb{E} \omega^4(z) - \bar{p}^2 = \mathbb{E} (\beta \bar{q})^2 \eta^4 \sigma^8 - \bar{p}^2. \end{aligned}$$

The last step missing is averaging over the η , by exploiting $\langle \eta^2 \rangle = 1$, $\langle \eta^4 \rangle = 3$. Finally, we have obviously $D(0) = E(0) = F(0) = 0$, because at $t = 0$ the two parties are independent.

Here, we are interested in finding where ergodicity becomes broken (the critical line), we start propagating $t \in 0 \rightarrow 1$ from the annealed region, where $\bar{q} \equiv 0$ and $\bar{p} \equiv 0$.

It is immediate to check that, for the only terms that we need to consider,

A, D, G (the other being strictly zero on the whole $t \in [0, 1]$), the starting points are $A(0) = 1, D(0) = 0, G(0) = (1 - \beta)^{-2}$ and their evolution is ruled by

$$\dot{A} = 2AD, \quad (37)$$

$$\dot{D} = AG + D^2, \quad (38)$$

$$\dot{G} = 2GD. \quad (39)$$

So we need to solve the system above. The first step is noticing that

$$d_t \log A = \frac{\dot{A}}{A} = 2D = \frac{\dot{G}}{G} = d_t \log G,$$

as $d(A/G)/dt = 0$, and $A(0)/G(0) = (1 - \beta)^2$, we obtain immediately the coupled behavior of the self-correlations:

$$A(t) = G(t)(1 - \beta)^2. \quad (40)$$

We now reduced to consider the system

$$\dot{D} = (1 - \beta)^2 G^2 + D^2, \quad (41)$$

$$\dot{G} = 2GD. \quad (42)$$

Let us call $[D + (1 - \beta)G] = Y$ such that summing (41) and (42) we get the differential equation

$$\dot{Y}(t) = Y^2(t) \Rightarrow Y(t) = \frac{Y_0}{1 - tY_0},$$

by which, as $Y_0 = (1 - \beta)^{-1}$, we get

$$D(t = \sqrt{\alpha}\beta) + (1 - \beta)G(t = \sqrt{\alpha}\beta) = \frac{1}{1 - \beta(1 + \sqrt{\alpha})}, \quad (43)$$

i.e. there is a regular behavior up to $\beta_c = 1/(1 + \sqrt{\alpha})$.

Now, starting from eq.(43), we have to solve separately for $G(t)$ and for $D(t)$.

Let us at first notice that

$$\dot{G}(t) = 2G(t)\left(Y(t) - (1 - \beta)G(t)\right), \quad (44)$$

by which, dividing both the sides by G^2 and considering $Z = G^{-1}$, we get

$$-\dot{Z}(t) - 2Y(t)Z(t) + 2(1 - \beta) = 0, \quad (45)$$

namely an ordinary first order differential equation for $Z(t)$.

We solve it by posing $Z(t) = W(t) \exp \left(- 2 \int_0^t Y(t') dt' \right)$, with $Z_0 = W_0$ fixing the auxiliary function $W(t)$ as

$$\int_0^t Y(t') dt' = \log \left(\frac{1 - \beta}{1 - \beta - t} \right).$$

We can obtain in a few algebraic steps the function $Z(t)$ and consequently, remembering that $G(t) = Z^{-1}(t)$ we get

$$G(t) = \frac{1}{2(1 - \beta)} \left(\frac{1}{1 - \beta - t} + \frac{1}{1 - \beta + t} \right) = \frac{1}{(1 - \beta)^2 - t^2}. \quad (46)$$

Now it is possible to insert eq.(46) into (43) which concludes the proof of the following

Theorem 2. *In the ergodic region the behavior of the overlap fluctuations is regular and described by the following equations:*

$$\langle \xi_{12}^2 \rangle = \frac{(1 - \beta)^2}{(1 - \beta)^2 - \beta^2 \alpha}, \quad (47)$$

$$\langle \xi_{12} \eta_{12} \rangle = \frac{\beta \sqrt{\alpha}}{(1 - \beta)^2 - \beta^2 \alpha}, \quad (48)$$

$$\langle \eta_{12}^2 \rangle = \frac{1}{(1 - \beta)^2 - \beta^2 \alpha}. \quad (49)$$

The ergodic region ends in the line

$$\beta_c = \frac{1}{1 + \sqrt{\alpha}}, \quad (50)$$

which is the critical line.

We stress that it turns out to be the same AGS-line of the standard neural network counterpart.

5 Conclusion and outlook

In this paper we achieved another step toward a general theory of neural networks whose statistical mechanics is not based on replica-trick.

We found the replica symmetric behavior of the analogical Hopfield model, its self-averaging equations for the order parameters and a complete quantitative picture of their fluctuations and correlations. The critical line defining

ergodicity breaking is found as well, in agreement with the standard AGS counterpart.

Furthermore the method paves the way for analytical investigation of general bipartite systems, which are assuming by themselves a very important role in applied statistical mechanics [14].

Despite these new results, fundamental enquiries are still open: apart the challenging thermodynamic limit, the retrieval phase (the response to an external stimulus) has not been discussed so far, neither the replica symmetry breaking scheme, which should be incorporated in the theory too.

We plan to report soon on these topics.

Acknowledgements

Support from MiUR (Italian Ministry of University and Research) and INFN (Italian Institute for Nuclear Physics) is gratefully acknowledged.

AB work is supported by the SmartLife Project (Ministry Decree 13/03/2007 n.368) which is acknowledged.

References

- [1] S. Albeverio, B. Tirozzi, B. Zegarlinski, *Rigorous results for the free energy in the Hopfield model*, Comm. Math. Phys. **150**, 337 (1992).
- [2] D.J. Amit, *Modeling brain function: The world of attractor neural network*, Cambridge University Press, (1992).
- [3] D.J. Amit, H. Gutfreund, H. Sompolinsky, *Spin Glass model of neural networks*, Phys. Rev. A **32**, 1007-1018, (1985).
- [4] D.J. Amit, H. Gutfreund, H. Sompolinsky, *Storing infinite numbers of patterns in a spin glass model of neural networks*, Phys. Rev. Lett. **55**, 1530-1533, (1985).
- [5] R. Albert, A. L. Barabasi, *Statistical mechanics of complex networks*, Rev. Mod. Phys. **74**, 47-97, (2002).
- [6] A. Barra, *Irreducible free energy expansion and overlap locking in mean field spin glasses*, J. Stat. Phys. **123**, 601-614, (2006).
- [7] A. Barra, F. Guerra, *About the ergodic regime in the analogical Hopfield neural networks: Moments of the partition function*, J. Math. Phys. **50**, 125217, (2008).

- [8] A. Barra, F. Guerra, *Constraints for the order parameters in analogical neural networks*, Percorsi d'Ateneo, S. Vitolo Ed., Salerno, (2008).
- [9] A. Bernacchia, D.J.Amit, *Impact of spatiotemporally correlated images on the structure of memory*, P.N.A.S. **104**, 3544, (2007).
- [10] A. Bovier, B. Niederhauser, *The spin-glass phase-transition in the Hopfield model with p-spin interactions*, Adv. Theor. Math. Phys. **5**, 1001 – 1046, (2001).
- [11] A. Bovier, A.C.D. van Enter, B. Niederhauser, *Stochastic symmetry-breaking in a Gaussian Hopfield-model*, J. Stat. Phys. **95**, 181-213, (1999).
- [12] A. Bovier, V. Gayrard, *An almost sure central limit theorem for the Hopfield model*, Markov Proc. Rel. Fields **3**, 151-173, (1997).
- [13] A. Bovier *Self-averaging in a class of generalized Hopfield models*, J. Phys. A **27**, 7069-7077, (1994).
- [14] P. Contucci, I. Gallo, *Bipartite Mean Field Spin Systems. Existence and Solution*, Math. Phys. Elec. Jou. **14**, 1-22, (2008).
- [15] A.C.C. Coolen, R. Kuehn, P. Sollich, *Theory of Neural Information Processing Systems*, Oxford University Press, (2005).
- [16] R.S. Ellis, *Large deviations and statistical mechanics*, Springer, New York, (1985).
- [17] A. Engel, C. Van den Broeck, *Statistical Mechanics of Learning*, Cambridge University Press, (2001).
- [18] F. Guerra, *An introduction to mean field spin glass theory: methods and results*, In: *Mathematical Statistical Physics*, A. Bovier et al. eds, 243 – 271, Elsevier, Oxford, Amsterdam, (2006).
- [19] F. Guerra, *Broken Replica Symmetry Bounds in the Mean Field Spin Glass Model*, Comm. Math. Phys. **233**, 1-12, (2003).
- [20] F. Guerra, *About the overlap distribution in mean field spin glass models*, Int. Jou. Mod. Phys. B **10**, 1675-1684, (1996).
- [21] F. Guerra, *Sum rules for the free energy in the mean field spin glass model*, in *Mathematical Physics in Mathematics and Physics: Quantum and Operator Algebraic Aspects*, Fields Institute Communications **30**, Amer. Math. Soc. (2001).

- [22] F. Guerra, F. L. Toninelli, *The Thermodynamic Limit in Mean Field Spin Glass Models*, Comm. Math. Phys. **230**, 71-79, (2002).
- [23] F. Guerra, F. L. Toninelli, *The infinite volume limit in generalized mean field disordered models*, Markov Processes and Rel. Fields, **9**, 195 – 207, (2003).
- [24] D.O. Hebb, *Organization of Behaviour*, Wiley, New York, (1949).
- [25] V. Honavar, L. Uhr, *Artificial Intelligence and Neural Networks: Steps Toward Principled Integration*, Elsevier, Boston, Academic Press, (1994).
- [26] J. Hertz, A. Krogh, R. Palmer, *Introduction to the theory of neural computation*, Santa Fe Institute Studies in the Sciences of Complexity, (1991).
- [27] J.J. Hopfield, *Neural networks and physical systems with emergent collective computational abilities*, P.N.A.S. **79**, 2554-2558, (1982).
- [28] M. Krein, A. Nudelman, *The Markov moment problem and extremal problems. Ideas and problems of P. L. Chebyshev and A. A. Markov and their further development*. Amer. Math. Soc. **50**, Providence, (1977).
- [29] M. Mézard, G. Parisi, M. A. Virasoro, *Spin glass theory and beyond*, World Scientific, Singapore, (1987).
- [30] L. Pastur, M. Shcherbina, *The absence of self-averaging of the order parameter in the Sherrington-Kirkpatrick model*, J. Stat. Phys. **62**, (1991).
- [31] L. Pastur, M. Scherbina, B. Tirozzi, *The replica symmetric solution of the Hopfield model without replica trick*, J. Stat. Phys. **74**, 1161-1183, (1994).
- [32] L. Pastur, M. Scherbina, B. Tirozzi, *On the replica symmetric equations for the Hopfield model*, J. Math. Phys. **40**, 3930-3947, (1999).
- [33] M. Talagrand, *Rigorous results for the Hopfield model with many patterns*, Prob. Theor. Relat. Fiel. **110**, 177-276, (1998).
- [34] M. Talagrand, *Exponential inequalities and convergence of moments in the replica-symmetric regime of the Hopfield model*, Ann. Prob. **38**, 1393-1469, (2000).
- [35] M. Talagrand, *Spin glasses: a challenge for mathematicians. Cavity and mean field models*, Springer Verlag, (2003).